

****TITLE****

*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***

****NAMES OF EDITORS****

Spectroscopy of BL Lac Objects: new redshifts and mis-identified sources.

N. Carangelo^{1,4}

¹ *Università di Milano-Bicocca, P.za della Scienza 3, 20126 Milano, I*
E-mail:nicoletta.carangelo@mib.infn.it

R. Falomo², J. Kotilainen³, A. Treves⁴ and M.-H. Ulrich⁵

² *Osservatorio Astronomico di Padova;* ³ *Tuorla Observatory;*

⁴ *Università dell'Insubria;* ⁵ *European Southern Observatory*

Abstract. We are carrying out a program of high signal to noise optical spectroscopy of BL Lacs with unknown or tentative redshift. Here we report some preliminary results. New redshifts are measured for PKS0754+100 ($z=0.266$) and 1ES0715-259 ($z=0.464$). From lineless spectra of PG1553+113 and PKS1722+119 we set a lower limit of $z>0.3$ for both sources. In two cases (UM493 and 1620+103) stellar spectra indicate a wrong classification.

1. Introduction and Observations

At variance with other classes of AGN BL Lac Objects are characterized by quasi featureless optical spectra. This often hinders the determination of the redshift (and therefore of the distance) of these sources. In the Véron Cetty and Véron catalog (2001) of AGN there are ~ 600 objects classified as BL Lacs and only for half of them the redshift is known. In order to contribute to the measurement of this fundamental parameter we are carrying out a systematic study of the optical spectra of objects with unknown or tentative redshift. We present here preliminary results from two campaigns at the European Southern Observatory (ESO) telescope performed in 2001 July and 2002 January.

The observations were gathered using the 3.6m telescope equipped with EFOSC2. Standard data reduction was performed using different packages in IRAF in order to obtain 1-dimensional wavelength and flux calibrated extracted spectra.

2. Results

2.1. New redshifts

PKS 0754+100 is a highly polarized and variable source that was discovered by Tapia et al. (1977) and classified as a blazar by Angel & Stockman (1980). A first tentative redshift proposed by Persic and Salucci (1986, $z=0.66$) was put in doubt after the detection of the host galaxy by Abraham et al. (1991) and by

Falomo (1996), because at this redshift the host would be extremely luminous ($M_R \simeq -25$).

Our spectrum (see fig. 1) shows clearly two emission lines at $\lambda=4717.2$ Å (EW=0.9 Å) and $\lambda=6340.4$ Å (EW=1.3 Å). Identification of these features with [OII] (3727 Å) and [OIII] (5007 Å) yields a redshift $z=0.266\pm0.001$. This is consistent with the tentative redshift proposed by Falomo and Ulrich (2000, $z\simeq0.28$). At $z=0.266$ the host galaxy luminosity becomes $M_R=-22.9$. We note that at the same redshift there is a companion galaxy, 13'' NE, corresponding to a projected distance of ~ 70 kpc (Pesce et al. 1995).

1ES 0715-259 belongs to the *Einstein* Slew Survey (Elvis et al. 1992) and was classified as a BL Lac Object by Perlman et al. (1996). They assumed that a VLA radio source could be identified with the corresponding Slew Survey source if it fell within 80'' of X-ray position. However in this case they reported an offset in the radio-X position of $>200''$. The VLA radio map of 1ES 0715-259 (Perlman et al. 1996) shows a clear FR II morphology. The MMT spectrum obtained by Perlman et al. (1996) is featureless which motivated the BL Lacs classification. The value of the ratio F_x/F_r ($\log F_x/F_r=-5.18$) places 1ES 0715-259 near the borderline between RBL and XBL objects. There are no data of polarimetry for this source. HST (Urry et al. 2000) clearly resolved 1ES0715-259 into a point source surrounded by a small, rather elongated host galaxy.

We took an optical spectrum of the counterpart of the radio source. This shows clearly strong emission lines at $\lambda=7114.5$ (EW=85.3 Å), $\lambda=7262.4$ Å (EW= 14.3 Å) and $\lambda=7332.0$ Å (EW=46 Å) (see figure 1). Identification respectively with the H_β , [OIII] 4959 Å and [OIII] 5007 Å yields a redshift $z=0.464\pm0.001$. The H_β luminosity ($L_{H_\beta}=2.5 \times 10^{43}$ erg s $^{-1}$) is typical of radio loud quasar. On the basis of our optical spectroscopy and the radio properties we believe that this source is not a BL Lac but a radio loud quasar. The mis-identification by Perlman et al. (1996) is likely due to the very poor quality of their optical spectrum.

2.2. Lineless objects

For some relatively bright objects ($V\sim 15-16$) despite the high signal to noise ($S/N\sim 200$) the spectra appear lineless. We report here two examples: PG1553+113 and PKS1722+119 (see figure 2).

PG 1553+113 is a very bright optically selected source (Green et al. 1986) classified as a BL Lacs on the basis of the featureless spectrum (Miller & Green 1983) and the large optical variability ($\Delta m \simeq 2$ mag, Miller et al. 1988). Falomo & Treves (1990) showed that the emission feature in the UV region identified as Lyman- α at $z=0.36$ by Miller & Green (1983) is spurious. Falomo & Treves (1990) did not find absorption features with an EW upper limit of 1 Å.

Our high signal to noise optical spectrum of the object confirms the absence of lines with a much lower of EW (~ 0.1 Å). The redshift of this source remains unknown. Assuming that it is hosted by a galaxy typical of BL Lacs ($M_R=-23.7$ $R_e=10$ kpc, Urry et al. 2000) from our spectrum we set a lower limit of $z>0.3$ consistent with the fact that the source is unresolved in HST images (Urry et




Figure 1. Left: The normalized spectrum of PKS0754+100. The two emission lines at $z=0.266$ are marked. Right: Spectrum of 1ES0715-259 showing clearly strong emission lines at $z=0.464$ of a typical QSO.




Figure 2. The normalized spectra of PG1553+113 (left) and PKS1722+119 (right). The spectra have $S/N=200$ and are lineless ($EW < 0.1 \text{ \AA}$).

al. 2000).

PKS1722+119 is a highly polarized X-ray selected BL Lac Object originally discovered by Uhuru. Griffith et al. (1989) claimed the observation of an absorption feature near 6000 Å which could be Na I (5892 Å) redshifted at $z=0.018$. This was disproved by Brissenden et al. (1990) who obtained a completely lineless spectrum, consistently with more recently results by Vèron Cètty and Vèron 1993 and by Falomo et al. 1993.

The spectrum presented here (see fig. 2) is lineless and we can set a lower limit of $z>0.3$. Also this source is unresolved by HST (Urry et al. 2000).

2.3. Mis-identified sources

From the spectra examined so far we found two cases UM493 and 1620+103 where we observe the typical stellar absorption lines of the Balmer series clearly indicating a wrong classification. Probably the mis-identifications (e.g. Vèron-Cètty & Vèron 2001, Hewitt & Burbidge 1993) were due to the poor informations on these sources, classification being based only on modest optical spectroscopy. Our high signal-to noise spectra clarify their nature and we can definitively classify these sources as stars of spectral type A.

References

- Abraham R.G., McHardy I.M., Crawford C.S. 1991, MNRAS, 252, 482
 Angel J.R.P. & Stockman H.S. 1980, ARA&A, 18, 321
 Brissenden R.J.V. et al. 1990, ApJ, 350, 578
 Elvis M. et al. 1992, ApJS, 80, 257
 Falomo R. & Treves A. 1990, PASP, 102, 1120
 Falomo R. et al. 1993, ApJ, 411, L63
 Falomo R. 1996, MNRAS, 283, 241
 Falomo R. & Ulrich M-H 2000, A&A, 357, 91
 Green R.F., Schmidt M. and Liebert J. 1986, ApJS, 61, 305
 Griffiths R.E. et al. 1989, MNRAS, 240, 33
 Hewitt A. & Burbidge G., 1993, ApJS, 87, 451
 Miller H.R. and Green R.F. 1983, *Bull. AAS*, 15, 957
 Miller H.R. et al. 1988, in proc. *IUE Symposium*, Greenbelt, ESA SP-281, p303
 Perlman E.S. and Stocke J.T. et al. 1996, ApJS, 104, 251
 Persic M. & Salucci P. 1986, *Struc. and Evol. of AGN*, ed Giuricin et al. , p657
 Pesce, J.E., Falomo, R. Treves, A., 1995, AJ, 110, 1554
 Tapia et al. 1977, ApJ, 215, 71
 Urry C.M. et al. 2000, ApJ, 532, 816
 Vèron-Cètty M.P. and Vèron P. 2001, A&A, 374, 92